

LA-UR-17-28706

Approved for public release; distribution is unlimited.

Title:	Least-squares elastic reverse-time migration of microseismic data using inverted moment-tensor sources
Author(s):	Chen, Yu Gao, Kai Huang, Lianjie
Intended for:	SEG 2017, 2017-09-25/2017-09-28 (Houston, Texas, United States)
Issued:	2017-09-28 (rev.1)

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Least-squares elastic reverse-time migration of microseismic data using inverted moment-tensor sources

Yu Chen, Kai Gao and Lianjie Huang

Geophysics Group, Los Alamos National Laboratory

Los Alamos, NM 87545, USA

Objective

Imaging fracture zones using least-squares reverse-time migration of microseismic data

Outline

- **Introduction**
- **Adaptive moment-tensor joint inversion**
- **Least-squares reverse-time migration**
- **Conclusions**

Outline

- **Introduction**
- Adaptive moment-tensor joint inversion
- Least-squares reverse-time migration
- Conclusions

Introduction


- Monitoring CO₂ reservoirs for long-term storage using induced microseismicity
- Using microseismic data
 - To directly image fracture zones using microseismic data, and
 - To imaging sedimentary layers surrounding microseismic clusters
- Conventional least-squares reverse-time migration using explosive or vector sources cannot match radiation patterns of microseismic data

Introduction


- We recently developed a new method for adaptive moment-tensor joint inversion of microseismic data acquired from a single-borehole geophone array.
- In this work:
 - We employ least-squares reverse-time migration of microseismic data using moment-tensor sources.
 - We apply the method to microseismic data acquired at the Aneth CO₂-Enhance Oil Recovery (EOR) field using single vertical borehole.

The Aneth CO₂-EOR field at Utah

- CO₂ injection from 2007 to 2009
- 23 levels of geophones within a vertical borehole spanning from 800 to 1700 m




(Soma and Rutledge, 2013)



Aneth CO₂-EOR field

- More than 3000 microseismic events detected from May 2008 to March 2009
 - 1266 events were relocated using direct, reflection and diffraction waves
- (Soma and Rutledge, 2013)




Outline

- Introduction
- **Adaptive moment-tensor joint inversion**
- Least-squares reverse-time migration
- Conclusions

Adaptive moment-tensor joint inversion method and its application to the Aneth CO₂-EOR field

- Uncertainty of moment-tensor inversion with limited azimuthal coverage
- Clustering events with similar seismic waveforms and radiation patterns
- Similar focal mechanisms in the adjacent areas (e.g. Dahm et al., 1999; Rutledge, 2004; Maxwell, 2014)
- Inverting the clustered events with the same focal mechanism





An adaptive joint inversion

- **Joint inversion**: events in a cluster are inverted using the same focal mechanism (strike, dip, rake, ISO and CLVD) but different source durations and moments
- **Adaptive inversion**: each event is further inverted based on the joint inversion result with a search range of $\pm 10^\circ$ for strike, dip and rake, ± 0.05 for ISO and CLVD

Synthetic test for joint inversion



50 events with the same focal: stk:220 dip:45 rake:0 iso:0.1 clvd:0.2

Same configuration as the Aneth field



Adaptive joint inversion with varying focal



50 events with varying focals: stk:220 dip:45 \pm (up to 10) rake:0 iso:0.1 clvd:0.2



Individual inversion


Left: full moment tensor

Right: double couple only




Joint inversion result

- Seven clusters based on similarities of microseismic waveforms and radiation pattern
- Consistent but slightly varying focal mechanisms of the seven clusters
- Large non-double couple component



Adaptive inversion result




Outline

- Introduction
- Adaptive moment-tensor joint inversion
- **Least-squares reverse-time migration**
- Conclusions

Least-squares reverse-time migration of microseismic data using inverted moment tensors

- Explosive source cannot make use of the strong S waves
- Explosive and vector source cannot match the radiation pattern of microseismic source




Moment-tensor source in finite difference

- Add the moment tensor as the stress in waveform modeling


$$\sigma_{ij} = \sigma_{ij} + \frac{\Delta t M_{ij}(t)}{V}, i = 1, 3, j = 1, 3$$

Synthetic test

True model




Smooth model



Up and Down going wavefield


Explosive sources

Moment-tensor sources





Right going wavefield

Explosive sources




Moment-tensor sources




Application to the Aneth field

- Selecting 38 microseismic events from the seven clusters
- Forming a 2D line
- Performing the imaging in a 3D model




Up and Down going wavefield

Explosive sources




Moment-tensor sources




Right going wavefield


Explosive sources




Moment-tensor sources



Explosive sources



Moment-tensor sources



Outline

- Introduction
- Adaptive moment-tensor joint inversion
- Least-squares reverse-time migration
- **Conclusions**

Conclusions

1. Developed an adaptive moment-tensor joint inversion method and applied to the Aneth field
 - The joint inversion inverts clustered events with the same focal mechanism
 - The adaptive inversion inverts each event in a search range around the joint inversion result
2. Developed a least-squares reverse-time migration algorithm for microseismic data
 - Image fracture zones around microseismic clusters
 - Image several sedimentary layers above and beneath the sources
 - Need to use moment-tensor sources to produce correct migration images

Acknowledgements

- The work was supported by the U.S. Department of Energy through contract DE-AC52-06NA25396 to Los Alamos National Laboratory.
- The computation was performed using the super-computer resources of LANL's Institutional Computing Program.

Thank you for your attention!